



Offshore winds from advanced synthetic aperture radar

Hasager, Charlotte Bay; Christiansen, Merete Bruun; Nielsen, Niels Morten

Publication date:
2008

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Hasager, C. B., Christiansen, M. B., & Nielsen, N. M. (2008). *Offshore winds from advanced synthetic aperture radar*. Abstract from 2007 European Offshore Wind Conference and Exhibition, Berlin, Germany.
http://www.risoe.dk/rispubl/art/2008_02_abstract.pdf

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

OFFSHORE WINDS FROM ADVANCED SYNTHETIC APERTURE RADAR

Charlotte Hasager (Risø national laboratory, DTU, Wind energy, Denmark)

Charlotte bay Hasager, Denmark (1) Merete bruun Christiansen, (1) Morten Nielsen, (1)

(1) risoe national laboratory, DTU

Onboard the European Space Agency (ESA) satellite Envisat the Advanced Synthetic Aperture Radar (ASAR) mainly observes in wide swath mode, i.e. with a swath of 400 km along the polar orbit track. Routine download of data and online calculation to offshore wind maps are done by Risoe using the ANSWRS software from the Johns Hopkins University, Applied Physics Laboratory, USA. Offshore wind resource statistics are investigated for the Danish Seas. The results are compared to model results and offshore observations.

The approach of offshore wind mapping using ASAR data from the Envisat satellite is to download the raw data in near-real-time, calibrate the data and apply a CMOD function using a priori wind direction as input. The a priori wind direction is taken from the NOGAPS atmospheric model. The relevance of using a series of offshore wind maps is to estimate the offshore wind climatology. It is innovative to use satellite information and it forms an alternative to offshore meteorological observation, however, only with accuracy adequate in pre-feasibility but not for wind farm siting as stand alone application. The results from the Danish Seas are presented by comparing to state of the art wind resource model results and offshore observations. The conclusion is that satellite ASAR provide a relevant regional offshore wind mapping capability for pre-feasibility. In addition, the data appear highly relevant for evaluation of offshore wind results from meso-scale modelling.